

November 2019

Sun	Mon	Tue	Wed	Thu	Fri	Sat
27	28	29	30	31	1	2
	CYCLE 1 Labs Davis Clinic Doxorubicin Infusion Ifosfamide Infusion Mesna @ __ PM	Urine Labs Mesna @ __ PM	Urine Labs Mesna @ __ PM	Dexamethasone 4mg 2x daily Ondansetron 8mg 2x daily Neulasta		
3	4	5	6	7	8	9
				Labs Davis Clinic		
10	11	12	13	14	15	16
17	18	19	20	21	22	23
	CYCLE 2 Labs Davis Clinic Doxorubicin Infusion Ifosfamide Infusion Mesna @ __ PM	Urine Labs Mesna @ __ PM	Urine Labs Mesna @ __ PM	Dexamethasone 4mg 2x daily Ondansetron 8mg 2x daily Neulasta		
24	25	26	27	28	29	30
				Labs Davis Clinic		
1	2	3	4	5	6	7

Nausea

Take Dexamethasone (Decadron) 4mg and Ondansetron (Zofran) 8mg twice daily for 3 days after infusion (see calendar).

For extra nausea, take below as needed:

- Level 1: Ondansetron (Zofran)
- Level 2: Prochlorperazine (Compazine)
- Level 3: Lorazepam (Ativan)

Port

You will have blood labs drawn on the first day of the cycle and approximately the 10th day. You can use Emla cream on your port to help numb it. Apply cream 40 minutes before and cover with plastic wrap.

Mesna

Mesna is a medication that will help protect your bladder from chemotherapy. You will take it at a different time everyday depending on what time your infusion started. Your nurse will tell you what time to take it.

Neulasta

Neulasta helps increase your white (infection-fighting) blood cells. On your third day of infusion, your nurse will put an on-body injector. It will inject the medication on the next day. It will beep before it starts injecting.

Mouth care

To decrease the risk of mouth sores, mix 1 tablespoon of baking soda and 1 tablespoon of salt with a glass of water. As you remember, rinse your mouth daily 3-4 times a day with this solution.

Fever

If your temperature exceeds 100.4 F, call the clinic. For pain, avoid using regular Acetaminophen (Tylenol), Ibuprofen (Advil, Motrin), Naproxen (Aleve), or Aspirin. These can mask signs of infection and impair platelet (blood clotting) cells.